

Antimicrobial Stewardship in Wound Care

7 Tips

Right patient, right treatment, right dose, right duration



1.

Regular hand disinfection and proper contact precautions are key in preventing wound infection



2.

Avoid prescribing antimicrobials when they are not indicated



3.

When antimicrobial therapy is indicated, prescribe an appropriate regimen



4.

Order therapy for the correct duration, at the optimal dose and by the appropriate route⁵



5.

Use an agent that has the least risk for adverse effects for the patient and the community^{6,7}



6.

Interdisciplinary collaboration helps inform and enforce AMS



7.

Every healthcare worker is responsible for being aware of local AMR issues and for undertaking AMS



Managing the Crisis of Antibiotic Resistance

The global prevalence for non-healing wounds (also referred to as chronic wounds, hard-to-heal wounds or complex wounds) with mixed aetiologies has been estimated at 2.21 per 1000 population.¹ The number of individuals developing non-healing wounds is increasing due to changing lifestyles² and an ageing population.³ A paper in *The Lancet* (2022) approximated that in 2019 there were 4.95 million global deaths associated with, and 1.27 million deaths directly attributable to, bacterial antimicrobial resistance (AMR).⁴ With the growing global problem of AMR, clinicians must use antibiotics wisely. This means following the principles of antimicrobial stewardship (AMS).

An antimicrobial strategy for non-healing wounds should include:

- Routinely determining if the wound is infected
- Surveillance programmes for wound infection
- Clear and achievable metrics
- Local policies to review the appropriateness of antimicrobial use
- An interdisciplinary team of specialists supervising AMS efforts
- Accessible multi-professional educational programmes
- Antimicrobial guardianship programmes
- Patient awareness campaigns




www.ewma.org



www.bsac.org.uk

"The most effective way to reduce the use of antimicrobials is to prevent infection."

A Concise Approach to Treating Potentially Infected Wounds



www.ewma.org



www.bsac.org.uk

The EWMA AMS programme is supported by an unrestricted educational grant from Convatec, Essity, Flen Health, Hydrofera Blue, Mölnlycke and PolyMem.

Contact: ewma@ewma.org

References

- Martinengo L, Olsson M, Bajpai R, Soljak M, Upton Z, Schmidchen A, et al. Prevalence of chronic wounds in the general population: systematic review and meta-analysis of observational studies. *Ann Epidemiol*. 2019 Jan;29:8-15.
- Gethin G, Touriani E, van Netten J, Sobotka L, Probst S. The impact of patient health and lifestyle factors on wound healing, Part 1: Stress, sleep, smoking, alcohol, common medications and illicit drug use. *J Wound Manag*. 2022;23(1 Suppl 1, pt 1):1-41.
- Gould L, Abadir P, Brem H, Carter M, Conner-Kerr T, Davidson J, et al. Chronic Wound Repair and Healing in Older Adults: Current Status and Future Research. *J Am Geriatr Soc*. 2015 Mar;63(3):427-38.
- Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, et al. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*. 2022 Feb;399(10325):629-55.
- International Wound Infection Institute (IWII). Wound infection in clinical practice. *Wounds International*. 2022.
- Lipsky BA. Diabetic foot infections: Current treatment and delaying the "post-antibiotic era". *Diabetes Metab Res Rev*. 2016 Jan;32 Suppl 1:246-53.
- Lipsky BA, Dryden M, Gottrup F, Nathwani D, Seaton RA, Stryja J. Antimicrobial stewardship in wound care: a Position Paper from the British Society for Antimicrobial Chemotherapy and European Wound Management Association. *J Antimicrob Chemother*. 2016 Nov;71(11):3026-35.
- Probst S, Apelqvist J, Bjarnsholt T, Lipsky BA, Ousey K, Peters EJG. Antimicrobials and Non-healing Wounds: An Update. *J Wound Management*, 2022;23(3 Sup1):S1-S33.